

A VECTOR FORM OF ALEKSANDROV'S THEOREM FOR NORMAL TOPOLOGICAL SPACES

SURJIT SINGH KHURANA

ABSTRACT. Let X be a Hausdorff normal topological space, E a quasi-complete locally convex space, $C(X)$ (resp. $C_b(X)$) the space of all (resp. all, bounded), scalar-valued continuous functions on X , and \mathcal{F} the algebra generated by the closed subset of X . The following form of Aleksandrov's theorem is proved: Suppose $\mu: C_b(X) \rightarrow E$ a weakly compact linear mapping. Then there exists a unique finitely additive, exhaustive measure $\nu: \mathcal{F} \rightarrow E$ such that

(i) ν is inner regular by closed sets and outer regular by open sets;

(ii) $\int f d\nu = \mu(f)$, $\forall f \in C_b(X)$.

When X is also countably paracompact some additional results are also proved.

1. INTRODUCTION AND NOTATION

In this paper \mathbb{R} stands for the set of real numbers, K will denote the field of real or complex numbers (we will call them scalars), X a Hausdorff normal topological space and E a quasi-complete locally convex space space over K with topology generated by an increasing family of semi-norms $\|\cdot\|_p$, $p \in P$; E' will denote the topological dual of E . For a $p \in P$, $V_p = \{x \in E : \|x\|_p \leq 1\}$; polars will be taken in the duality $\langle E, E' \rangle$. We denote by $C(X)$ the space of all K -valued continuous functions on X , and by $C_b(X)$ the bounded elements of $C(X)$. $Z \subset X$ will be called a zero-set if $Z = f^{-1}(0)$ for some $f \in C_b(X)$ and the complements of zero-sets will be called positive sets. The elements of the σ -algebra generated by zero-sets are called Baire sets and the elements of the σ -algebra generated by closed sets are called Borel sets; $\mathcal{B}(X)$ and $\mathcal{B}_0(X)$ are the classes of Borel and Baire subsets of X and $M_\sigma(X)$ denotes the class of all scalar-valued, countably additive Baire measures on X . For locally convex spaces, the notation and results of [11] will be used. For a vector space F , F^* will denote its algebraic dual. N will denote the set of natural numbers. For topological measure theory notations and results of ([5], [7], [12], [13], [14]) will be used. All locally convex spaces are assumed to be Hausdorff and over K . \tilde{X} will denote the Stone-Cech compactification of X .

Received: September 27, 2010. *Revised:* April 18, 2011.

2010 *Mathematics Subject Classification:* Primary: 46G10, 28C15, 28B05; Secondary: 28A30.

Key words and phrases: Measure representation of linear operators, Aleksandrov's theorem.

With sup-norm topology, $C_b(X)$ is a Banach space which is isometric isomomorphic to $C(\tilde{X})$. The dual of $C_b(X)$ is denoted by $M(X)$. For a function $f \in C_b(X)$, \tilde{f} will denote its unique continuous extension to \tilde{X} .

Now we come to vector-valued measures; integrability of scalar-valued functions is taken in the sense of [8]. If \mathcal{A} is a σ -algebra of subsets of a set Y , $\mu: \mathcal{A} \rightarrow E$ a countably additive vector measure and $p \in P$, we denote the p -semi-variation of μ by $\bar{\mu}_p$, $\bar{\mu}_p(A) = \sup\{|g \circ \mu|(A) : g \in V_p^0\}$ (here V_p^0 is the polar of V_p in the duality $\langle E, E' \rangle$) [8]; we consider the submeasure $\dot{\mu}_p: \mathcal{A} \rightarrow R^+$, $\dot{\mu}_p(A) = \sup\{\|\mu(B)\|_p : B \in \mathcal{A}, B \subset A\}$ ([2], [5]). It is easily verified that $\dot{\mu}_p$ is countably sub-additive [2] and $\dot{\mu}_p \leq \bar{\mu}_p \leq 4\dot{\mu}_p$. Also there is a control measure for $\bar{\mu}_p$; this will be denoted by λ_p ; this control measure can be chosen to be in the closed convex hull of $\{|g \circ \mu| : g \in V_p^0\}$, with norm topology on measures ([8], p. 20, proof of Theorem 1). This control measure also has the properties that:

(i) $|f \circ \mu| \ll \lambda_p$ for every f in E' with $\|f\|_p \leq 1$;

(note that $\|f\|_p = \sup\{|f(x)| : x \in V_p\}$);

(ii) if $\lambda_p(A) = 0$ then $\bar{\mu}_p(A) = 0$;

(iii) $\lim_{\lambda_p(A) \rightarrow 0} \bar{\mu}_p(A) = 0$;

(iv) $\lambda_p \leq \bar{\mu}_p$.

We also have the result that if $f: Y \rightarrow K$ is measurable function, $B \in \mathcal{A}$ and $|f| \leq c$ on B , then $\left\| \int_B f d\mu \right\|_p \leq c\bar{\mu}_p(B)$.

$L^1(\mu)$ will denote the space of μ -integrable functions [8]. For any $f \in L^1(\mu)$, we take $\bar{\mu}_p(f) = \sup\{|g \circ \mu|(|f|) : g \in V_p^0\}$ ([8], Lemma 2, p. 23).

If \mathcal{F} is an algebra of subsets of a set Y and $\mu: \mathcal{F} \rightarrow E$ a finitely additive measure then μ will be called exhaustive if for any disjoint sequence $\{A_n\} \subset \mathcal{F}$, we have $\mu(A_n) \rightarrow 0$ ([2]); exhaustive measures are called strongly bounded measures in [1]; for quasi-complete E , a finitely additive μ is exhaustive iff $\mu(\mathcal{F})$ is relatively weakly compact in E - for Banach spaces, it is proved in [1] and it easily extends to quasi-complete locally convex spaces.

2. ALEKSANDROV'S THEOREM

In this section, we extend the celebrated Aleksandrov representation theorem, for Hausdorff normal topological spaces, to the vector-valued measures. In scalar case, in a simple form, this theorem says:

Suppose X is a Hausdorff normal topological space, \mathcal{F} the algebra generated by the closed subset of X and $\mu: C_b(X) \rightarrow K$ a continuous linear mapping. Then there exist a unique, finitely additive measure $\nu: \mathcal{F} \rightarrow K$ such that:

(i) ν is inner regular by closed sets and outer regular by open sets;

(ii) $\int f d\nu = \mu(f), \forall f \in C_b(X)$. ([3], p.262; [13]). Note $C_b(X)$ is contained in the uniform closure of \mathcal{F} -simple functions on X in the space of all bounded functions on X and so each $f \in C_b(X)$ is ν -integrable.

We state and prove the following extension. Our proof is obtained by the regularity properties of the corresponding regular Borel measure on \tilde{X} and is very different from the given in [3] and other known proofs.

We start with a crucial lemma for normal topological spaces:

Lemma 2.1. *Suppose A and B are two closed sets in a Hausdorff normal space X . Then $\overline{A \cap B} = \overline{A} \cap \overline{B}$ (for a subset $C \subset X, \overline{C}$ denotes the closure of C in \tilde{X}).*

Proof. By normality, the result is true if $A \cap B = \emptyset$. So we assume $A \cap B \neq \emptyset$. Assume that $a \in \overline{A \cap B} \setminus \overline{A \cap B}$. So there is a closed neighborhood Q of a , in \tilde{X} , such that $Q \cap (A \cap B) = \emptyset$. Putting $P_1 = A \cap Q$ and $P_2 = B \cap Q$, we get $P_1 \cap P_2 = \emptyset$. Since X is normal, $\overline{P_1} \cap \overline{P_2} = \emptyset$. But $a \in \overline{P_1}$ and also $a \in \overline{P_2}$; this is a contradiction. \square

Now we come to the main theorem

Theorem 2.1. *Suppose X is a Hausdorff normal topological space and a weakly compact linear mapping $\mu: C_b(X) \rightarrow E$. Then there exist a unique finitely additive, exhaustive measure $\nu: \mathcal{F} \rightarrow E$ such that:*

- (i) ν is inner regular by closed sets and outer regular by open sets;
- (ii) $\int f d\nu = \mu(f), \forall f \in C_b(X)$.

Proof. Considering $\tilde{\mu}: C(\tilde{X}) \rightarrow E$, we shall get an E -valued regular Borel measure $\tilde{\mu}: \mathcal{B}(\tilde{X}) \rightarrow E$ ([7]) (we denote the measure $\tilde{\mu}$ by the same notation as operator $\tilde{\mu}$ because the operator $\tilde{\mu}$ uniquely extends to all $\tilde{\mu}$ -integrable functions). If A is a subset of X or \tilde{X} , \overline{A} will denote the closure of A in \tilde{X} . Fix $p \in P$.

We prove this theorem in several steps.

I. Let $\overline{\mathcal{C}} = \{\overline{A} : A \text{ a closed set in } X\}$. Then for every $Q \in \overline{\mathcal{C}}$ and $c > 0$, there exists $W \in \overline{\mathcal{C}}$, such that $W \subset \tilde{X} \setminus Q$ and $\tilde{\mu}_p(\tilde{X} \setminus Q \setminus W) < c$.

Proof. By regularity of $\tilde{\mu}_p$, there is a positive set $V \subset \tilde{X} \setminus Q$ having the property that $\tilde{\mu}_p(\tilde{X} \setminus Q \setminus V) < \frac{c}{2}$ (a positive set in the complement of a zero-set). Take a $g \in C(\tilde{X}), 0 \leq g \leq 1$, such that $V = g^{-1}(0, 1]$. Denote by

$$V_n = \left\{ x \in \tilde{X} : g(x) > \frac{1}{n} \right\}$$

and

$$Z_n = \left\{ x \in \tilde{X} : g(x) \geq \frac{1}{n} \right\}.$$

Now,

$$Z_{n+1} \supset \overline{(Z_{n+1} \cap X)} \supset \overline{(V_n \cap X)} \supset V_n$$

(note X is dense in \tilde{X}). By choosing n sufficiently large, without loss generality, we can assume that $\tilde{\mu}_p(V \setminus V_n) < \frac{c}{2}$. Taking $W = \overline{(Z_{n+1} \cap X)}$, we get the result. \square

II. Let \mathcal{A} be the algebra, in \tilde{X} , generated by $\overline{\mathcal{C}}$ and denote, by \mathcal{A}_0 , the elements of \mathcal{A} which have the property that these elements and their complements are inner regular by the elements of $\overline{\mathcal{C}}$. Then $\mathcal{A}_0 = \mathcal{A}$.

Proof. We use Step I and Lemma 2.1 to prove it. By Step I, $\mathcal{A}_0 \supset \overline{\mathcal{C}}$. By definition, \mathcal{A}_0 is closed under complements. Also, using Lemma 2.1, it is a routine verification that if A and B are in \mathcal{A}_0 then $A \cap B$ and $A \cup B$ are also in \mathcal{A}_0 . This proves the result. \square

III. It is a simple verification that $\mathcal{A} \cap X \supset \mathcal{F}$. Also if $A \in \mathcal{A}$ and $A \cap X = \emptyset$, then $\tilde{\mu}_p(A) = 0$. To prove this, take any $\overline{C} \in \overline{\mathcal{C}}$, C being a closed set in X , such that $\overline{C} \subset A$. This means C is empty and so $\tilde{\mu}_p(A) = 0$.

Now we can define a $\nu: \mathcal{F} \rightarrow E$, $\nu(B) = \tilde{\mu}(A)$, A being any element in \mathcal{A} with $B = A \cap X$; it is a trivial verification that ν is well-defined, is finitely additive and it is inner regular by closed sets in X and outer regular by open sets in X (this means for a $p \in P$, $F \in \mathcal{F}$ and $c > 0$, there is, in X , a closed set C and an open set V , $C \subset F$ and $V \supset F$, such that for any $B_1 \in \mathcal{F}$, $B_1 \subset F \setminus C$ and any $B_2 \in \mathcal{F}$, $B_2 \subset V \setminus F$, we have $\|\nu(B_i)\|_p < c$ for $i = 1, 2$). We also have $\nu(C) = \tilde{\mu}(\overline{C})$, for any closed $C \subset X$. Since $\nu(\mathcal{F})$ is relatively weakly compact in E , ν is exhaustive (\equiv strongly additive) ([1], Corollary 3, p. 28; this is proved for Banach space E but easily extends to quasi-complete locally convex space E). Also for any $B \in \mathcal{F}$, $\tilde{\nu}_p(B) \leq \tilde{\mu}_p(A)$, where A is any element in \mathcal{A} such that $B = A \cap X$.

IV. For any $f \in C_b(X)$, $\mu(f) = \int f d\nu$.

Proof. Assume $\tilde{\mu}_p(X) \leq 1$. Fix $c > 0$ and take an $f \in C_b(X)$, $0 \leq f \leq 1$. Then there is a non-negative, \mathcal{F} -simple function $\sum_{i=1}^n a_i \chi_{B_i}$ such that B_i 's are mutually

disjoint, their union is X and $\left| f - \sum_{i=1}^n a_i \chi_{B_i} \right| < c$ on X .

Take mutually disjoint $\{A_i\} \subset \mathcal{A}$ such that $B_i = A_i \cap X$ for every i . Also consider mutually disjoint closed sets $\{C_i\} \subset X$, such that $\tilde{\mu}_p(A_i \setminus \overline{C_i}) < \frac{c}{n}$, for each i . Now

$$\begin{aligned} \left\| \int f d\nu - \sum a_i \nu(C_i) \right\|_p &\leq \left\| \int f d\nu - \sum a_i \nu(B_i) \right\|_p + \left\| \sum a_i \nu(B_i \setminus C_i) \right\|_p \\ &\leq c + \left\| \sum a_i \tilde{\mu}(A_i \setminus \overline{C_i}) \right\|_p \leq c + n \cdot \frac{c}{n} = 2c. \end{aligned}$$

Also $\left| f - \sum_{i=1}^n a_i \chi_{B_i} \right| \leq c$ implies that

$$\left| \tilde{f} - \sum_{i=1}^n a_i \chi_{\overline{C_i}} \right| \leq c,$$

on $\cup(\overline{C_i})$ (note $\overline{C_i}$ are also mutually disjoint by Lemma 2.1).

Therefore,

$$\begin{aligned} \left\| \int \tilde{f} d\tilde{\mu} - \sum a_i \nu(C_i) \right\|_p &= \left\| \int \tilde{f} d\tilde{\mu} - \sum a_i \tilde{\mu}(\overline{C_i}) \right\|_p \\ &\leq c + \left\| \sum a_i \tilde{\mu}(A_i \setminus \overline{C_i}) \right\|_p \\ &\leq c + n \cdot \frac{c}{n} = 2c. \end{aligned}$$

This proves that $\mu(f) = \nu(f)$. □

V. Uniqueness.

Proof. Let $\nu: \mathcal{F} \rightarrow E$ be a finitely additive regular (inner regular by closed sets in X and outer regular by open sets in X) measure, having relatively weakly compact range, such that $\int f d\nu = 0, \forall f \in C_b(X)$. This means ν is exhaustive and so $\bar{\nu}_p(X) < \infty, \forall p \in P$. If $\nu \neq 0$, there is a $p \in P$, a closed set $Z \subset X$, and a $c > 0$ such that $\|\nu(Z)\|_p = 2c$. Take a open set $U \supset Z$ such that $\bar{\nu}_p(U \setminus Z) < c$. Take an $f \in C_b(X), 0 \leq f \leq 1, f(Z) = 1, f(X \setminus U) = 0$. We get

$$0 = \int f d\nu = \int_Z f d\nu + \int_{U \setminus Z} f d\nu.$$

This means $\nu(Z) = - \int_{U \setminus Z} f d\nu$ and so $2c \leq 1, \bar{\nu}_p(U \setminus Z) < c$. This contradiction proves the uniqueness. □

These steps prove the result. □

Now we make the additional assumption that X is also countably paracompact. In this case we prove that any Baire measure can be uniquely extended to regular Borel measure.

Theorem 2.2. *Suppose X is a Hausdorff normal and countably paracompact topological space and $\mu: \mathcal{B}_0(X) \rightarrow E$ a Baire measure. Then it has a unique extension to a countably additive Borel measure $\mu: \mathcal{B}(X) \rightarrow E$ which is inner regular by closed sets and outer regular by open sets.*

Proof. By ([7], Thorem 2), we get a unique linear, continuous, and weakly compact mapping $\mu: C_b(X) \rightarrow E$ which is continuous in the strict topology β_σ (which is weaker than norm topology) on $C_b(X)$. We fix a $p \in P$ and use the notations of Theorem 2.1. As in Theorem 2.1, we get an E -valued regular Borel measure $\tilde{\mu}: \mathcal{B}(\tilde{X}) \rightarrow E$.

We do the proof in several steps:

I. Let $\{C_n\}$ be a decreasing sequence of closed subsets of X such that $\cap C_n = C$. Then $\tilde{\mu}_p(\cap(\overline{C_n}) \setminus \overline{C}) = 0$.

Proof. We first consider the case when $C = \emptyset$. In this case $\{V_n = C'_n : 1 \leq n < \infty\}$ is an opening covering X . Take an open neighborhood finite refinement covering and a partition of unity $\{g_\alpha : \alpha \in I\}$ subordinated to this refinement. Let $f_n = \sum_{\alpha \in I_n} g_\alpha$

where $I_n = \{\alpha : \text{support of } g_\alpha \subset V_n\}$. We get $f_n \uparrow 1$ and so $Z_n = f_n^{-1}(0) \supset C_n$ and $\cap Z_n = \emptyset$. Note Z_n are zero-sets. Since μ is a Baire measure, we have $0 = \tilde{\mu}_p(\cap(\overline{Z_n}))$. Since $\cap(\overline{C_n}) \subset \cap(\overline{Z_n})$, we get $\tilde{\mu}_p(\cap(\overline{C_n})) = 0$. Now we consider the case when $C \neq \emptyset$. Fix a $c > 0$. By Theorem 2.1 (I), there is a closed set $Q \subset X \setminus C$ such that $\tilde{\mu}_p(\tilde{X} \setminus \overline{C} \setminus \overline{Q}) < c$. Since $\cap(C_n \cap Q) = \emptyset$, by using which is just proved above and Lemma 2.1, we get $\tilde{\mu}_p(\cap(\overline{C_n} \cap \overline{Q})) = 0$.

Now

$$\begin{aligned} \tilde{\mu}_p(\overline{C_n} \setminus \overline{C}) &= \tilde{\mu}_p((\overline{C_n} \cap \overline{Q}) \cup (\overline{C_n} \setminus \overline{Q} \setminus \overline{C})) \\ &\leq \tilde{\mu}_p(\overline{C_n} \cap \overline{Q}) + \tilde{\mu}_p(\tilde{X} \setminus \overline{Q} \setminus \overline{C}). \end{aligned}$$

Taking limits, we get $\tilde{\mu}_p(\cap(\overline{C_n}) \setminus \overline{C}) < c$. This proves the result. \square

II. In the notations of Theorem 2.1(I, II), let, this time, \mathcal{A} be the σ -algebra, in \tilde{X} , generated by $\overline{\mathcal{C}}$ and denote, by \mathcal{A}_0 , the elements of \mathcal{A} which have the property that these elements and their complements are inner regular by the elements of $\overline{\mathcal{C}}$. Then $\mathcal{A}_0 = \mathcal{A}$.

Proof. As done in Theorem 2.1(II), $\mathcal{A}_0 \supset \overline{\mathcal{C}}$ and it is an algebra. Take a sequence $\{A_n\} \subset \mathcal{A}_0$. Fix a $c > 0$ and take a sequence $\{\overline{C_n}\} \subset \overline{\mathcal{C}}$ such that $\tilde{\mu}_p(A_n \setminus \overline{C_n}) < \frac{c}{4^n} \forall n$. From this it easily follows that $\tilde{\mu}_p(\cup A_n \setminus (\bigcup_{1 \leq i \leq m} \overline{C_i})) < c$ for some m and

$\tilde{\mu}_p(\cap A_n \setminus \cap \overline{C_n}) < c$. Putting $C = \bigcup_{1 \leq i \leq m} C_i$, we get $\tilde{\mu}_p(\cup A_n \setminus \overline{C}) < c$. Let $D_n =$

$\bigcap_{1 \leq i \leq n} C_i$; then $D_n \downarrow$ and $\cap C_n = \cap D_n$ and so $\cap \overline{C_n} = \cap \overline{D_n}$. Put $D = \cap D_n$. By I,

$\tilde{\mu}_p(\cap(\overline{D_n}) \setminus \overline{D}) = 0$ and we also have $\tilde{\mu}_p(\cap A_n \setminus \cap \overline{C_n}) < c$. This gives

$$\begin{aligned} \tilde{\mu}_p(\cap A_n \setminus \overline{D}) &= \tilde{\mu}_p((\cap A_n \setminus \cap \overline{D_n}) \cup (\cap \overline{D_n} \setminus \overline{D})) = \tilde{\mu}_p((\cap A_n \setminus \cap \overline{C_n}) \cup (\cap \overline{D_n} \setminus \overline{D})) \\ &\leq \tilde{\mu}_p(\cap A_n \setminus \cap \overline{C_n}) + \tilde{\mu}_p(\cap \overline{D_n} \setminus \overline{D}) \leq c, \end{aligned}$$

(using again I).

This proves $\mathcal{A} = \mathcal{A}_0$. This proves the result. \square

III. $\mathcal{A} \cap X \supset \mathcal{B}(X)$

Proof. From $\mathcal{A} \supset \overline{\mathcal{C}}$, it follow that $\mathcal{A} \cap X$ contains all closed subset of X . Using this with the easily verified fact that $\mathcal{A} \cap X$ is a σ -algebra, the result follows. \square

IV. For any $A \in \mathcal{A}$ with $A \cap X = \emptyset$, $\tilde{\mu}_p(A) = 0$

Proof. By II, A is inner regular by the elements of $\overline{\mathcal{C}}$. Since $A \cap X = \emptyset$, the result follows. \square

Now we can define a $\mu: \mathcal{B}(X) \rightarrow E$. Take any $B \in \mathcal{B}(X)$ and select any $\tilde{B} \in \mathcal{A}$ such that $\tilde{B} \cap X = B$; define $\mu(B) = \tilde{\mu}(\tilde{B})$; it is a trivial verification that μ is well-defined, is countably additive and it is inner regular by closed sets in X and outer regular by open sets in X . \square

Remark 2.1. For scalar valued measure, this result is prove in [10].

3. REPRESENTATION THEOREM FOR $C(X)$, WITH X COMPLETELY REGULAR

In this section we assume that $K = \mathbb{R}$. A subset $B \subset C(X)$ will be called order-bounded if there are elements f and g in $C(X)$ such that $f \leq b \leq g, \forall b \in B$. It is well-known that a linear map $\mu: C(X) \rightarrow \mathbb{R}$, which maps order-bounded sets into bounded sets, gives a unique $\nu \in M_\sigma(X)$ such that $C(X) \subset L^1(\nu)$ and $\mu(f) = \int f d\nu$ ([4]; [14], Theorem 23).

We will extend to the vector case.

Theorem 3.1. *Suppose X is a Hausdorff normal and countably paracompact topological space and $\mu: C(X) \rightarrow E$ be a linear map such that order-bounded subsets are mapped into relatively weakly compact subsets of E . Then:*

(i) *there is a unique Borel measure $\nu: \mathcal{B}(X) \rightarrow E$, which is inner regular by closed subsets of X and outer regular by open subsets of X , such that $C(X) \subset L^1(\nu)$ and $\mu(f) = \int f d\nu$;*

(ii) *for every $p \in P$, there is compact $C \subset vX$ (the real-compactification of X), depending on p , such that $\tilde{\nu}_p(\tilde{X} \setminus C) = 0$ ([4]).*

Proof. By [7], Theorem 7, p. 695, there is $\nu: \mathcal{B}_0(X) \rightarrow E$, a unique Baire measure, satisfying all the conditions of the above theorem, except Borel extension. But the regular Borel extension follows form Theorem 2.2. \square

Acknowledgment. We are very thankful to the referee for pointing out some typographical errors and also making some very useful suggestions which have improved the paper. Also our thanks to Professor M. Rajagopalan for providing a simple proof of Lemma 2.1.

REFERENCES

- [1] J. Diestel and J. J. Uhl: *Vector Measures*, Amer. Math. Soc. Surveys, Vol. 15, Amer. Math. Soc., 1977.
- [2] L. Drewnowski: *Topological rings of sets, continuous set functions, integration I, II*, Bull. Acad. Polon. Sci. Ser. Math. Astronom. Phys., **20**(1972), 269-286.
- [3] N. Dunford and J. T. Schwartz: *Linear Operators*, Interscience Publishers, 1988.
- [4] E. Hewitt: *Linear functionals on spaces of continuous functions*, Fund. Math., **37**(1950), 161-189.
- [5] S. S. Khurana: *Topologies on spaces of continuous vector-valued functions*, Trans. Amer. Math. Soc., **241**(1978), 195-211.
- [6] S. S. Khurana: *Topologies on spaces of continuous vector-valued functions II*, Math. Ann., **234**(1978), 159-166.
- [7] S. S. Khurana: *Vector measures on Topological spaces*, Georg. Math. J., **14**(2007), 687-698.
- [8] I. Klivanek and G. Knowles: *Vector Measures and Control Systems*, North-Holland/American Elsevier, 1975.
- [9] D. R. Lewis: *Integration with respect to vector measures*, Pac. J. Math., **33**(1970), 157-165.
- [10] J. Marik: *The Baire and Borel measure*, Czech. Math. J., **7(82)**(1957), 248-253.
- [11] H. H. Schaeffer: *Topological Vector Spaces*, Springer Verlag, 1986.
- [12] F. D. Sentiiles: *Bounded continuous functions on completely regular spaces*, Trans. Amer. Math. Soc., **168**(1972), 311-336.
- [13] R. F. Wheeler: *Survey of Baire measures and strict topologies*, Expos. Math., **2**(1983), 97-190.
- [14] V. S. Varadarajan: *Measures on topological spaces*, Amer. Math. Soc. Transl., **48**(1965), No. 2, 161-228.

The University of Iowa
Department of Mathematics
Iowa City, Iowa, 52242, USA
E-mail address: surjit-khurana@uiowa.edu